

Pericardial infection caused by *Brevibacterium casei*

In response to the article describing two cases of catheter-related bacteraemia associated with *Brevibacterium casei* in the May 2004 issue of *Clinical Microbiology and Infection* [1], we wish to report an unusual case of *B. casei* infection that was encountered recently at our institution.

A 78-year-old man with a history of cancer was admitted with fatigue, shortness of breath, loss of appetite, and intermittent chest pain. The patient was afebrile, but hypotensive (blood pressure, 68/39 mmHg), tachycardiac (heart rate, 124/min), tachypenic (respiratory rate, 28/min) and hypoxic (oxygenation saturation, 88% on room air). On examination, there was an irregular heart rhythm, but without murmurs or gallop or peripheral signs of endocarditis. Initial laboratory values included a white blood cell count of $11\,100/\text{mm}^3$ (84% segmented neutrophils, 6% lymphocytes, 8% monocytes, 1% eosinophils and 1% basophils), haemoglobin 8.3 g/dL, and platelets $202\,000/\text{mm}^3$. Blood and urine cultures were negative. A trans-thoracic echocardiogram revealed a moderate-to-large pericardial effusion with right ventricle collapse. An emergent pericardiocentesis drained 570 mL of thick, bloody fluid. Pericardial fluid analysis revealed $140\,000$ red blood cells/ mm^3 , 208 white blood cells/ mm^3 (73% neutrophils, 27% lymphocytes), protein 4.8 g/dL, glucose 160 mg/dL, amylase 13 U/L and lactic dehydrogenase 4219 U/dL. Cytopathological analysis of the pericardial fluid showed acute and chronic inflammatory cells with reactive mesothelial cells. Culture of the pericardial fluid was positive for *B. casei*, and treatment with intravenous vancomycin was started.

Antimicrobial sensitivity testing was performed in conventional Gram-positive MicroScan (Sacramento, CA, USA) panels via broth microdilution, and was interpreted according to the breakpoints recommended by Funke *et al.* [2]. The organism was sensitive to ceftriaxone ($\text{MIC} \leq 4$ mg/L), ciprofloxacin ($\text{MIC} \leq 1$ mg/L), erythromycin ($\text{MIC} \leq 0.5$ mg/L), rifampicin ($\text{MIC} \leq 1$ mg/L), tetracycline ($\text{MIC} \leq 4$ mg/L) and vancomycin ($\text{MIC} \leq 2$ mg/L), but was resistant to ampicillin ($\text{MIC} 4$ mg/L), amoxycillin-clavulanic acid ($\text{MIC} > 4/2$ mg/L), clindamycin ($\text{MIC} > 2$ mg/L) and penicillin ($\text{MIC} 2$ mg/L).

The patient's condition improved initially, but a gastrointestinal bleed and atrial fibrillation, with rapid ventricular response, developed after a further 11 days, and the patient subsequently died.

The genus *Brevibacterium* was first described in the 1950s [3], but *Brevibacterium* spp. have been implicated only rarely as human pathogens [1,3–15]. *Brevibacterium* spp. are Gram-positive coryneform bacilli that are found frequently in dairy products. Some species (e.g., *B. casei* and *Brevibacterium epidermidis*) may be part of the normal human skin flora [3,16]. To date, *Brevibacterium* spp. have been identified in clinical specimens, but *B. casei* appears to be the most common [17]. Fourteen case reports of *Brevibacterium* spp. infections can be found in the literature [1,4–15]. Manifestations have included bacteraemia, with or without associated central venous catheter infection ($n = 10$), peritonitis ($n = 2$), endocarditis ($n = 1$) and sternal osteomyelitis ($n = 1$). Most clinical *Brevibacterium* isolates have been reported to be susceptible to β -lactams, ciprofloxacin and vancomycin. To our knowledge, the present case is the first report of pericardial infection caused by *Brevibacterium* spp. These organisms appear to be opportunistic pathogens, with most reported cases involving patients with an underlying malignancy or immunodeficiency disease. The possibility of *Brevibacterium* infection should be considered in the context of pericarditis among immunocompromised patients.

J. P. Cannon*, S. L. Spadoni, S. Pesh-Iman and
S. Johnson

Pharmacy Service and Medicine Service,
Hines VA Hospital,
Fifth Avenue,
and Roosevelt Road,
Hines, IL 60141, USA

*E-mail: Joan.Cannon2@med.va.gov

REFERENCES

1. Beukinga I, Rodriguez-Villalobos H, Deplano A, Jacobs F, Struelens MJ. Management of long-term catheter-related *Brevibacterium* bacteremia. *Clin Microbiol Infect* 2004; **10**: 465–467.
2. Funke G, Punter V, Von Graevenitz A. Antimicrobial susceptibility patterns of some recently established coryneform bacteria. *Antimicrob Agents Chemother* 1996; **40**: 2874–2878.

3. Gruner E, Steigerwalt AG, Hollis DG *et al.* Human infections caused by *Brevibacterium casei*, formerly CDC groups B-1 and B-3. *J Clin Microbiol* 1994; **32**: 1511–1518.
4. Wauters G, Van Bosterhaut B, Avesani V *et al.* Peritonitis due to *Brevibacterium otitidis* in a patient undergoing continuous ambulatory peritoneal dialysis. *J Clin Microbiol* 2000; **38**: 4292–4293.
5. Brazzola P, Zbinden R, Rudin C, Schaad UB, Heininger U. *Brevibacterium casei* sepsis in an 18-year-old female with AIDS. *J Clin Microbiol* 2000; **38**: 3513–3514.
6. Dass KN, Smith MA, Gill VJ, Glodstein SA, Lucey DR. *Brevibacterium* endocarditis: a first report. *Clin Infect Dis* 2002; **35**: e20–21.
7. Ogunc D, Gultekin M, Colak D *et al.* Bacteremia caused by *Brevibacterium* species in a patient with chronic lymphocytic leukemia. *Haematologia* 2002; **32**: 151–153.
8. Janda WM, Tipirneni P, Novak RM. *Brevibacterium casei* bacteremia and line sepsis in a patient with AIDS. *J Infect* 2003; **46**: 61–64.
9. Lina B, Carlotti A, Lesaint V, Devaux Y, Freney J, Fleurette J. Persistent bacteremia due to *Brevibacterium* species in an immunocompromised patient. *Clin Infect Dis* 1994; **18**: 487–488.
10. Antoniou S, Dimitriadis A, Polydorou F, Malaka E. *Brevibacterium iodum* peritonitis associated with acute urticaria in a CAPD patient. *Perit Dial Int* 1997; **17**: 614–615.
11. McCaughey C, Damani NN. Central venous line infection caused by *Brevibacterium epidermidis*. *J Infect* 1991; **23**: 211–212.
12. Neumeister B, Mandel T, Gruner E, Pfyffer GE. *Brevibacterium* species as cause of osteomyelitis in a neonate. *Infection* 1993; **21**: 177–178.
13. Reinert RR, Schnitzler N, Haase G *et al.* Recurrent bacteremia due to *Brevibacterium casei* in an immunocompromised patient. *Eur J Clin Microbiol Infect Dis* 1995; **14**: 1082–1085.
14. Kaukoranta-Tolvanen SS, Sivonen A, Kostiala AA, Hormila P, Vaara M. Bacteremia caused by *Brevibacterium* species in an immunocompromised patient. *Eur J Clin Microbiol Infect Dis* 1995; **14**: 801–804.
15. Castagnola E, Conte M, Venzano P *et al.* Broviac catheter-related bacteremias due to unusual pathogens in children with cancer: case reports with literature review. *J Infect* 1997; **34**: 215–218.
16. Koneman EW, Allen SD, Janda WM, Schreckenberger PC, Winn WC, eds. The aerobic Gram-positive bacilli. In: *Diagnostic microbiology*, 5th edn. Hagerstown, MD: Lippincott, 1997; 651–688.
17. Funke G, Carlotti A. Differentiation of *Brevibacterium* spp. encountered in clinical specimens. *J Clin Microbiol* 1994; **32**: 1729–1732.